

## Phytochemical Investigation And Pharmacological Evaluation Of Medicinal Plant Extracts

Ms. Shagufta A. Farooqui<sup>1\*</sup>, Dr. S.S Patil<sup>2</sup>, Ms. S. A Tekale<sup>3</sup>, Mr V.N Gunjkar<sup>4</sup>

<sup>1\*</sup>Research Scholar, School of Pharmacy SRTMUN, Maharashtra

<sup>2</sup>Principal Maharashtra College of Pharmacy, Nilanga, Maharashtra

<sup>3</sup>Research Scholar, Nanded Pharmacy college, Nanded, Maharashtra.

<sup>4</sup>Research Scholar, Nanded Pharmacy college, Nanded, Maharashtra.

### ABSTRACT:

The objective of the study was to investigate the Phytochemicals of Ethyl acetate and Methanol extracts of *Annona reticulata* L. stem bark.

The present study reports physicochemical characterization, antioxidant of stem bark extracts from *Annona reticulata* L. plant collected from local region of Nanded, Maharashtra, India, and authenticated by botanist. Different physical parameters like ash values, extractive value, Loss on drying, etc. were evaluated for powdered drug. The extracts were obtained from Soxhlet method by using Petroleum ether, Ethyl acetate and methanol as solvents for extraction and subjected for preliminary phytochemical evaluation and antioxidant studies.

Total phenolic and flavonoids content were also analyzed. The presence of primary and secondary metabolites such as carbohydrate, proteins, alkaloids, phenolic compounds, was confirmed through preliminary phytochemical analysis. DPPH free radical scavenging assays showed antioxidant activities with increase in concentration of Ethyl acetate and Methanol stem bark extracts.

**Keywords:** *Annona reticulata* L. Ethyl acetate and methanol extracts, Anti-oxidant activity.

### I. INTRODUCTION:

Plants have been recognized as one of the sources of the medicinal products which are helpful to mankind. A different plant shows different pharmacological activities. Extracts obtained from various parts of plant possess various medicinal properties. They are also used for cosmetic preparations as an herbal product shows lesser side effects as compared to the synthetic products. The repetitive or regular consumption of the synthetic drugs results in addiction or toxicity thus it is safer to use herbal medicines than the synthetic medicines.

Many developing countries use the plant-based products for their needs. Our country India has developed the traditional system known as Ayurvedic system which is based on herbal products. Ayurveda have developed uncountable herbal products which helps the people according to their needs with the safe environment. A medicinal plant typically contains mixture of different phytochemical which may improve quality of life.

*Annona reticulata* Linn. (Bullock's heart) is one of the traditionally important plants used for the treatment of various diseases. It belongs to family Annonaceae. Numerous phytoconstituents have been identified from different parts of *A. reticulata*. Stem bark contains tannins, alkaloid and phenolic compounds.

Leaves contain wide range of chemicals like alkaloids, amino acids, carbohydrates, steroids, flavonoids, proteins, tannins, glycosides and phenolic. The root has been identified for the content of acetogenin, alkaloid, carbohydrates, proteins, flavonoids, tannins.

Traditionally the plant has been employed for the treatment of epilepsy, dysentery, cardiac problem, parasite and worm infestations, constipation, hemorrhage, bacterial infection, dysuria, fever, ulcer and as insecticide. Bark is a powerful astringent and used as a tonic whereas leaves used for helminthiasis treatment. *Annona reticulata* Linn has been used to the treatment of pain and inflammation. Also, it reduces hyperthyroidism, and hyperglycaemia.

### Morphology:

Rampal is a small tree with a rounded crown and trunk of 25-35 cm thick its height ranges from 5-10 meter the leaves are alternate oblong and lanceolate, 10-20 cm long and 2-5cm wide. Flowers are fragrant collateral with narrow petals of 2-3cm long the flowers never fully open. The fruits are nearly heart shaped 8-16cm wide. The skin of fruits may be yellow or brownish when ripe with brownish red blush there is thick, cream- white layer of custard like flesh under the skin with moderately juicy segments. Actual seed counts 55, 60 and 76. A pointed central core is attached to the thick stem of the fruit.

### Geographical distribution:

It is specially found in tropical regions of India. It is also found in west indies and central and south America, Taiwan, Bangladesh, Australia and West Africa.



Image No.1: *Annona reticulata* L. Plant

Table 1 Scientific classification

Scientific classification	
Kingdom	Plantae
Order	Magnoliids
Family	Annonaceae
Genus	Annona
Species	<i>Annona reticulata</i>

Table 2 Botanical, Common Names

Botanical, common and vernacular names	
Botanical name	<i>Annona reticulata</i> Linn.
Common name	Netted Custard apple
English	Bullock's heart,
Corazon Portuguese	Frutoda-Condessa
Indonesian	Buah nona
India	Ramphal

### Medicinal uses:

The plant has been using to treatment of various diseases like epilepsy, parasite, cardiac problem, worm infection, hemorrhage, dysuria, fever, ulcer.

The **barks** of this plant have astringent property that's why this plant used as a tonic.

The **leaves** of *Annona reticulata* L. used for the helminthiases. also, the plant has been used as an anti-inflammatory in wound healing, antianxiety, anti-stress, spasmolytic effects.


Leaf and stem extracts show positive inotropic, and spasmolytic effect.

The **fruit** is used as an ant dysenteric and anthelmintic. *Annona reticulata* Linn has been used to the treatment of pain and inflammation. Also, it reduces hyperthyroidism, and hyperglycaemia.

## II. MATERIALS AND METHODS:

### Collection, identification and authentication of plant material

The fresh stem bark of *Annona reticulata* L. were collected from local region and were authenticated by **Dr. Shrirang S. Bodke**, Associate professor and Head of Botany and Horticulture, Yashwant Mahavidyalaya, Nanded. Authentication of plant *Annona reticulata* L. was done by specimen No. **H-2**. Collection, authentication, identification, processing and storage had been done according to standard procedure for the plant material.




**Image No.2: Herbarium sheet and Authentication letter**

#### A. Pharmacognostic Evaluation of Plant Material:

Colour	Dark brown
Taste	Slightly aromatic
Shape	Cylindrical
Odour	Characteristic

#### Microscopic characteristics

Transverse section of stem bark of *Annona reticulata* L. The thin section of stem was cut with the sharp blade and stained with phloroglucinol and dil. HCl (1:1) and presence or absence of following characteristic was observed.




**Image No.3: TS of plant Stain with Phloroglucinol + HCL**

**Image No.4: TS of plant Without stain**

#### Powdered characteristics of stem bark

Some amount of powdered drug stained with phloroglucinol and HCL (1:1) it gives pink color to fibers which was observed under microscope with 40X lens.



**Image No.5: Powdered characteristics Stain with Phloroglucinol + HCL**

**Image No.6: Powdered characteristics without stain**

### Processing of crude drug.

The fresh stem barks of plant *Annona reticulata* L. were subjected to shade drying and further crushed to coarse powder, and then the powder passed through mesh No. 14 and stored in air tight container for further use.



Image No.7: Processing of crude drug

### B. Physicochemical Evaluation and Phytochemical Screening:

#### i. Total Ash value

Porcelain dish or a tarred silica crucible ignited and weighed. About 2 g of powder was weighed into dish/crucible. the dish Support on pipe-clay triangle placed on ring of stand. Heated using burner, with flame about 2 cm high and supporting the dish about 7 cm above the flame, heated till vapours almost cease was evolved, after that lowered the dish and heated more strongly until all the carbon is burnt off cooled in to desiccator. Ash was weighed and the total ash value was calculated the with reference to the air-dried sample of the crude drug (Khandelwal k. 2002).

#### ii. Acid Insoluble Ash

Proceeded as per the steps was mentioned in the procedure for determination of total ash value of a crude drug. After that using 25 ml of dilute hydrochloric acid, ash was washed from the dish used for total ash into 100 ml beaker. The mere gauze was placed was placed over a Bunsen burner and boiled for 5 min. It is filtered with an 'ash less' filter paper, and residue was washed twice. Crucible ignited on in the flame, cooled and weighed. The filter-paper and residue were put together into crucible, heated gently until vapours cease was evolved and then more strongly until all carbon has been removed. Cooled in to desiccator the residue was weighed and acid-insoluble ash calculated with the crude drug with reference to the air-dried sample of the crude drug. (Khandelwal K. 2002)

#### iii. Water Soluble Ash

Proceeded as per the steps was mentioned in the procedure for determination of total ash value of a crude drug. After that using 25 ml of water, ash was washed from the dish used for total ash into 100 ml beaker. The mere gauze was placed over a Bunsen burner and boiled for 5 min. It is filtered with an 'ash less' filter paper, and residue was washed twice. Crucible ignited on in the flame, cooled and weighed. The filter-paper and residue were put together into crucible, heated gently until vapours cease was evolved and then more strongly until all carbon has been removed. Cooled in to desiccator. The residue was weighed and water-soluble ash calculated with the crude drug with reference to the air-dried sample of the crude drug. (Khandelwal K. 2002)



Image No.8: Burning of ash



#### iv. Determination of Loss on Drying (LOD)

Weighed about 2 g of the powdered drug into a flat and thin porcelain dish. Dried in the oven at 105°C, until two consecutive weighing do not differ by more than 0.5 mg. Cooled in desiccator and weighed. The loss in weight is usually recorded as moisture. (Khandelwal K. 2002)

#### v. Extractive Value

4 gm of air-dried drug was macerated with 100 ml of solvents like pet ether, chloroform, acetone, methanol etc. and the specified strength in a closed flask for 24 hours. It was frequently Shaked during the first 6 hours and allowed to stand for 24 hours. Thereafter, it was filtered rapidly without loss of solvents, and dried in petri dish, and weighed. The percentage of extractive value was calculated (Khandelwal K. 2002).



Image No.09: Extractive value stand for 24 hr. & Evaporation

#### C. Extraction of Plant Material:

##### i. Selection of Solvent

On the basis extractive value and nature of phytochemical present in drug and literature review solvents were selected for the extraction of the Stem bark of *Annona reticulata* L. like Petroleum ether, Ethyl acetate & Methanol.

##### ii. Selection of Extraction Method

Study of literature survey revealed that most of the chemical constituents of the plant extract are heat stable and most of the researchers selected continuous hot extraction method for plant extraction, **Soxhlet extractor** is very essential with less time usage and with high efficiency, solvent penetrates faster to the plant and it is most convenient method. on this basis the **Soxhlet extraction method** was selected for extraction of stem bark powder of *Annona reticulata* L.




Image No.10: Soxhlet Extraction Ethyl acetate and Methanol Extraction

### Preparation of extracts

Three extracts of Stem bark of *Annona reticulata* L. was prepared

- i. Petroleum ether extracted by continuous hot extraction method
- ii. Ethyl acetate extracted by continuous hot extraction method
- iii. Methanol extracted by continuous hot extraction method

The extract obtained and the dried mass was weighed and recorded. The percentage of yield was calculated.

$$(\%) \text{ yield} = \frac{\text{Wt. of extract}}{\text{Wt. of powdered drug}} \times 100$$

#### ➤ Preparation of Petroleum ether extract

Dried powdered plant was successfully extracted with petroleum ether by Soxhlet extractor apparatus according to the standard method till colorless solution was observed in siphon tube. 270 gm of the powdered plant and 1500 ml petroleum ether was used for extraction. After completion of extraction extract was cooled & evaporated by using Superfit Rotary evaporator. The extract was stored in air tight container. % yield of extract was calculated.

#### ➤ Preparation of Ethyl acetate extract

Dried powdered plant was successfully extracted with Ethyl acetate by Soxhlet extractor apparatus according to the standard method till colorless solution was observed in siphon tube. 260 gm of the powdered plant and 1500 ml Ethyl acetate was used for extraction. After completion of extraction extract was cooled & evaporated by using Superfit Rotary evaporator. The extract was stored in air tight container. % yield of extract was calculated.

#### ➤ Preparation of Methanol extract

Dried powdered plant was successfully extracted with methanol by Soxhlet extractor apparatus according to the standard method till colorless solution was observed in siphon tube. 250 gm of the powdered plant and 1500 ml methanol was used for extraction. After completion of extraction extract was cooled & evaporated by using Superfit Rotary evaporator. The extract was stored in air tight container. % yield of extract was calculated.

### D. Phytochemical Qualitative Analysis of Extract

#### i. Chemical tests

The extracts obtained by successive extraction were subjected to qualitative tests for the identification of various secondary metabolites such as carbohydrates, proteins, tannins, steroids, flavonoids, alkaloids and glycosides. Phytochemical examinations were carried out for all the extracts as per standard methods.

#### ➤ Test for carbohydrate:

##### a) Molisch test:

The aqueous extract treated with few drops of alpha-naphthol solution in alcohol, shake and add conc.  $\text{H}_2\text{SO}_4$  from sides of the test tube. Violet ring is formed at the junction of two liquids.

##### b) Fehling's test:

1 ml Fehling's A and 1 ml Fehling's B solution was mixed, boiled for one minute. Equal volume of test solution added. Heated on boiling water bath for 5-10 min. First yellow then brick red ppt is observed.

##### c) Benedict's test:

Mix equal volume benedict's reagent and test solution in test tube. Heat in boiling water bath for 5min solution appears green, yellow or red depending on amount of reducing sugar present in the test solution.

#### ➤ Test for proteins:

##### a) Biuret test:

Mix 3 ml T.S. Add 4 % NaOH and few drops of 1%  $\text{CuSO}_4$  solution. Violet or pink colour appears.

##### b) Million's test:

Mix 3 ml T.S with 5 ml Million's reagent. White ppt. turns brick red or the ppt. dissolves giving red coloured solution.

##### c) Test for Protein containing sulphur:

Mix 5 ml T.S. with 2ml 40% NaOH and 2 drops 10 % lead acetate solution. Boil mixture. Solution turns black or brownish due to  $\text{PbS}$  formation.

#### ➤ Test for amino acids:

##### a) Ninhydrin test (general test):

Heat 3 ml T.S. and 3 drops 5% Ninhydrin solution in boiling water bath for 10 min. purple or bluish colour appears.

#### ➤ Test for steroids:

##### a) Salkowski reaction:

To 2ml of extract at 2 ml chloroform and 2 ml conc.  $\text{H}_2\text{SO}_4$ . Shake well chloroform layer appears red and acid layer shows greenish yellow fluorescence.

**b) Liebermann-Burchard reaction:**

Mix 2ml extract with chloroform add 1-2 ml acetic anhydride and 2 drops conc.  $H_2SO_4$  from side of test tube. First red then blue finally green colour appears.

➤ **Test for glycosides:**

**a) Legal's test:**

To aqueous or alcoholic extract, add 1 ml pyridine and 1 ml sodium nitroprusside. Pink to red colour appears.

**b) Keller-killiani:**

To 2 ml extract, add glacial acetic acid, one drop 5%  $FeCl_3$  and conc.  $H_2SO_4$ . Reddish brown colour appears at junction of the two liquid layers and upper layer appears bluish green.

➤ **Test for saponins:**

**a) Foam test:**

shake the drug extract or dry powder vigorously with water. Persistent stable foam observed.

➤ **Test for flavonoids:**

**a) Shinoda test:**

To dry powder or extract added 5 ml 95% methanol few drops conc.  $H_2SO_4$  and 0.5 gm. magnesium turnings. Orange, pink, red to purple colour appears. Added t-butyl alcohol before adding the acid to avoid accidents from a Violent reaction and to dissolve coloured compounds in to the upper phase by using zinc instead of magnesium, only flavones give a deep red to magenta colour while flavanones and flavonols give weak pink to magenta colour or no colour is observed.

**b) Sulphuric acid test:**

On addition of sulphuric acid (66 or 80%) flavones and flavonols dissolve into it and give a deep yellow solution. Chalcones and aurones give red, bluish solutions. Flavanones give orange to red colours.

➤ **Test for tannins and phenolic compounds**

To 2-3 ml of aqueous or alcoholic extract, add few drops of following reagents:

**a) Lead acetate test:**

bark extract was taken and 0.5 ml of 1% lead acetate solution was added and the formation of white ppt. observed.

**b) 5% Ferric chloride solution:**

Deep blue-black colour is observed.

**c) Dilute iodine solution:**

Transient red colour.

**d) Bromine water:**

Decolouration of bromine water.

➤ **Test for alkaloids:**

The small portion of solvent free extract/ fraction was stirred separately with a few drops of dilute HCL, filtrated and used to test the presence of alkaloids:

**a) Mayer's test:**

2-3 ml filtrate with few drops of Mayer's reagent gives ppt.

**b) Hager's test:**

2-3 ml filtrate with Hager's reagent gives yellow ppt. indicates presence of alkaloids.

**c) Wagner's test:** 2-3 ml filtrate with few drops of Wagner's reagent gives reddish brown ppt.

**ii. Thin Layer Chromatography**

➤ **Introduction**

Thin layer chromatography is a method of analysis in which the stationary phase, a finely divided solid, is spread as a thin layer on a rigid supporting plate and the mobile phase, a liquid, is allowed to migrate across the surface of the plate by capillary action by gravity or pressure. TLC separation takes place in the open layer with each component having the same total migration time but different migration distance. Numerous fixed sorbents have been used, including Silica gel, Cellulose, Polyamide, Alumina, Ion exchange and chemically bonded silica gel. Mobile phase consists of a single solvent or a mixture of solvents. The stationary phase of the TLC is prepared using various techniques such as pouring, dipping and spraying.

The prepared plates are allowed for setting (air drying). This is done to avoid cracks on the surface of adsorbent. After setting the plates are activated by keeping in an oven at 100 to 200°C for one hour. Activation of TLC plates is nothing but removing water/moisture and other substances from the surface of any adsorbent, by heating at temperature around 1100°C so that adsorbent activity is retained. TLC studies were carried out using various extracts to confirm the presence of different Phytoconstituents in the extracts.

➤ **Analysis**

In TLC qualitative analysis of the unknown compound is done by comparing the  $R_f$  values. As solutes never travel the

full length of the stationary phase in TLC all the Rf value depends on the amount of the stationary phase, the humidity, layer thickness, solvent quality, saturation of chamber, development distance, temperature, amount of substance added, and the presence of impurities.

$$Rf = \frac{\text{Distance Travelled by solute}}{\text{Distance travelled by Solvent}}$$

Rf = Retention factor

#### ➤ Development of TLC profile of the extracts

All the extracts of selected plant material were subjected to TLC studies using various solvent systems to determine the presence of various Phytoconstituents. The Rf of observed compounds were noted for all extracts. The characteristic of the various chemical constituents in each extract under UV light and after derivatization with suitable reagents was recorded.

Preliminary phytochemical screening revealed the presence of flavonoids, Tannins, glycoside, amino acid. Compounds of varying polarity in the extracts well separated using various solvent systems on TLC. Rf value of the separated compounds were recorded. (Cai, Li et.al 2014)

### E. Phytochemical Quantitative Analysis of Extracts:

#### i. Total Phenolic content

##### ➤ Principle:

The Total phenolic content of the extract was determined by the Folin–Ciocalteu method. This test is based on oxidation of phenolic groups with phosphomolybdate and phosphotungstate. After oxidation a green-blue complex formed which measure at 666 nm.

##### ➤ Chemical required:

Folin-Ciocalteu (1:10 diluted with water)

Sodium carbonate (7%)

Extract (50 µg/ml)

##### ➤ Procedure

The Total phenolic content of the extract was determined by the Folin–Ciocalteu method. Extracts (50 µg/ml) were made up to 3 ml with distilled water, mixed thoroughly with 0.5 ml of Folin-Ciocalteu reagent for 3 min, followed by the addition of 2 ml of 7% (w/v) Sodium Carbonate. Then final volume made up to 10 ml.

The mixture was allowed to stand for a further 60 min in the dark, and absorbance was measured at 666 nm. The total phenolic content was calculated from the calibration curve of gallic acid (2, 4, 6, 8, 10 µg/ml) and by using equation obtained from standard calibration curve ( $r^2 = 0.9919$ ) and the results were expressed as mg per gallic acid equivalent per gram dry weight. (Subba, Bimala et.al 2016)

#### ii. Total Flavonoid content

##### ➤ Principle:

This method is based on the nitration of any aromatic ring bearing a catechol group with its 3-4 position unsubstituted or not sterically blocked. After addition of  $AlCl_3$  a yellow solution of complex was formed which then turned immediately to red after addition of NaOH and the value of absorbance is measured at 510 nm.

##### ➤ Chemical required:

Aluminum trichloride (10%)

Distilled water

Sodium hydroxide (1molar)

Sodium nitrate 5%

Extract solution (50 µg/mL)

##### ➤ Procedure

The total flavonoid content of crude extract was determined by the aluminum chloride colorimetric method. 1ml extract solution (50 µg/mL) was mixed with 4ml distilled water and 0.3 ml of 5%  $NaNO_2$  solution and 0.3 ml of 10%  $AlCl_3$  solution was added after 5 min add 2 ml /1molar NaOH, and the mixture was allowed to stand for 6 min, and the final volume of the mixture was brought to 10 ml with distilled water.

The mixture was kept for 30 min, and absorbance was measured at 510 nm against blank (without standard). quercetin was used as standard. The total flavonoid content was calculated from a calibration curve, (quercetin 2,4,6,8,10 µg/ml) and by using equation obtained from standard calibration curve ( $r^2=0.9911$ ) result was expressed as mg quercetin



equivalent per gram dry weight. (Subba, Bimala et.al 2016)

## F. Pharmacological Screening of Plant Extracts

### ➤ In-vitro studies

In-vitro study of Stem bark of *Annona reticulata* L. is done in which, following study was carried out which includes,

#### i. In-vitro Anti-oxidant Activity

An antioxidant is a molecule capable to inhibiting the oxidation of other molecules. Oxidation is a chemical reaction that transfers electrons or hydrogen from a substance to an oxidizing agent. Oxidation reactions can produce free radicals. these radicals can start chain reactions. When the chain reaction occurs in a cell, it can cause damage or death to the cell. Antioxidants terminate these chain reactions by removing free radical intermediates, and inhibit other oxidation reactions.

#### A. DPPH (2, 2-dipheny-1-picrylhydrazyl) radical scavenging activity.

##### ➤ Principle

DPPH assay method is based on the reduction of Methanolic solution of colored free radical DPPH by free radical scavenger. DPPH composed of stable free radical molecules. The DPPH radical contains an odd electron, which is responsible for the absorbance at 517 nm and also for a visible deep purple color. The procedure involves measurement of decrease in absorbance of DPPH at its absorption maxima of 517 nm, which is proportional to concentration of free radical scavenger added to DPPH reagent solution. The activity is expressed as effective concentration EC<sub>50</sub>.

##### ➤ Reagents Required

DPPH (0.1 mM)

Methanol

Quercetin (standard)

Extract solution

##### ➤ Preparation of samples and standard solutions

Accurately weighed 10 mg of Ethyl acetate and Ethanol extracts and the standard Quercetin dissolved in methanol. These solutions were serially diluted with methanol to obtain the lower dilutions.

##### ➤ Procedure:

The ability of *Annona reticulata* L. stems bark to scavenge DPPH radicals was measured according to the method as previously described with some modifications. Initially, absorbance of DPPH solution (400mg in 10 ml methanol 0.1 mM) was read at decreasing wavelength 513 nm. Various concentrations of extracts 20,40,60,80,100 µg/ml) were mixed with 0.4 ml of 0.1 mM DPPH radical solution in methanol and made up to final volume of 10 ml with methanol. A similar solution without samples was applied as a control. The mixture was shaken vigorously and incubated in the dark for 1 hr. at room temperature. Quercetin was used as a standard. The absorbance was measured at decreasing wavelength 513 nm by using uv-visible spectrophotometer. The percent radical scavenging activity of tested samples was expressed by using following formula.

$$\% \text{ Scavenging Activity} = \frac{\text{Control absorbance} - \text{Test absorbance}}{\text{Control absorbance}} \times 100$$

## III OBSERVATIONS AND RESULTS

### A. Physical Evaluation of Plant Material

#### i.Total ash value

Table 2 Total ash value of *Annona reticulata* L.

Parameters	(Z)Wt. of crucible+ ash(gm)	(X)Wt. of empty crucible(gm)	(Y)Wt. of drug (gm)	% of ash =Z-X÷Y×100
Total ash value	26.55	26.40	2	7.5%
Acid insoluble ash	44.06	44.05	2	0.5%
Water soluble ash	26.47	26.40	2	2%

#### ii.Determination of Loss on Drying (LOD)

Loss on drying can be calculated using following formula.

$$\text{LOD} = \frac{(W2 - W3)}{(W2 - W1)} \times 100$$

W1 = wt. of empty dish = 67.10

W2 = wt. of dish + drug = 68.69

W3 = wt. of drug after drying = 68.40

$$\text{LOD} = \frac{(68.69 - 68.40)}{(68.69 - 67.10)} \times 100$$

**LOD = 18%**

### iii.Extractive value

**Table 3 Extractive value of *Annona reticulata* L.**

Sr. No.	Solvents	(a)Wt. of petri dish with lead(gm)	(b)Wt. of petri dish with extraction (gm)	Wt. of extract (gm)	Percent of extract (%) =b-a÷4×100
1.	Pet ether	87.29	87.55	0.26	6.5%
2.	Chloroform	92.28	92.37	0.09	2.25%
3.	Acetone	92.23	92.27	0.04	1%
4.	Ethyl acetate	81.68	81.74	0.06	1.5%
5.	Ethanol	88.11	88.27	0.16	4%
6.	Methanol	90.39	90.61	0.22	5.5%
7.	Water	86.30	86.62	0.32	8%

### iv.Extraction of plant material

**Table 4 % of Yield of *Annona reticulata* L. Stem bark extracts**

Sr. No.	Drug taken (g)	Solvent used (ml)	Consistency	Colour of extract	Yield (g)	(%) Yield
1.	270	1500 (Pet Ether)	Sticky	Dark Green	1.26	0.46
2.	260	1500 (Ethyl Acetate)	Sticky	Dark brown	3.62	1.41
3.	240	1500 (Methanol)	Sticky	Dark brown	2.56	1.06



**Image No.11: AR-ME &AR-EAE**

### B. Phytochemical screening of extracts


**Table 5 Chemical tests of all three Stem bark extracts *Annona reticulata* L.**

Chemical tests	Pet ether	Ethyl acetate	Methanol
<b>Test for Carbohydrate</b>			
Molisch test	-	+	+
Fehling's test	-	+	+
Benedict's test	-	+	-
Barford's test	-	-	-
<b>Test for Proteins</b>			

<b>Biuret test</b>	+	+	-
<b>Millon's test</b>	-	+	-
<b>Test for protein S.</b>	-	+	+
<b>Test for Amino acid</b>			
<b>Ninhydrin test (general test)</b>	-	-	-
<b>Test Steroid</b>			
<b>Salkowski test</b>	-	+	+
<b>Liebermann Burchard reaction</b>	-	-	-
<b>Test for Glycosides</b>			
<b>Legal's test</b>	-	+	-
<b>Keller-killiani test</b>	-	-	-
<b>Test for Saponin Glycosides</b>			
<b>Foam test</b>	-	+	+
<b>Test for Flavonoids</b>			
<b>Shinoda test</b>	-	+	+
<b>Sulphuric acid test</b>	-	+	+
<b>Test for Tannins and phenolic compound</b>			
<b>Lead acetate test</b>	-	+	+
<b>5% Ferric chloride test</b>	-	+	+
<b>Dil. iodine solution</b>	-	-	-
<b>Bromine water</b>	-	-	+
<b>Test for Alkaloids</b>			
<b>Dragendroff's test</b>	-	-	-
<b>Wagner's test</b>	-	+	+
<b>Mayer's test</b>	-	+	+
<b>Hager's test</b>	-	+	+

**(+) Present, (-) Absent**

Above observation table shows the presence of phytoconstituents in the *Annona reticulata* L. Stem bark extracts. It reveals that most of the chemical constituents present Ethyl acetate and Methanol extract. It contains carbohydrates, alkaloids, steroids, flavonoids, phenols, proteins, amino acids only contain in Ethyl acetate and Methanol extract not present in Petroleum ether extract.






Image No.12: Chemical Tests of *Annona reticulata* L. stem bark extracts

#### Thin Layer Chromatography

Thin Layer Chromatography of *Annona reticulata* L. stem bark extracts.



Image No.13: TLC of AR-PE



Image No.14: TLC of AR-ME





Image No.15: TLC of AR-EAE

# Rf values of *Annona reticulata* L. stem bark extracts

**Table 6 Rf values of All three extracts of *Annona reticulata* L.**

Sr. No.	Extracts	Solvent System	Proportion	Spraying reagent	Rf Value	Colour	Chemical constituent
1.	AR-PE	Toluene: Ethyl Acetate: Methanol	8:1:1	10% H <sub>2</sub> SO <sub>4</sub>	0.52	Purple	Triterpenoids
2.					0.77	Yellow	Flavonoids
3.					0.87	Purple	Triterpenoids
4.	AR-ME	Chloroform: Ethyl acetate: Methanol	7:2:0.5		0.23	Light Brown	Flavonoids
5.					0.38	Brown	Alkaloid
6.					0.61	Pink	Diterpenes
7.					0.90	Faint Green	Chlorophyll
8.	AR-EAE	n-hexane: Ethyl acetate: Methanol	5:4:0.5		0.29	Faint Brown	Alkaloid
9.					0.51	Brown	Flavonoids
10.					0.63	Faint pink	Diterpenes
11.					0.80	Brown	Alkaloid
12.				0.92	Green	Chlorophyll	


## C. Phytochemical quantitative analysis of extracts

### i. Total Phenolic content

**Table 7 Absorbance of Gallic acid at different concentrations for determination of total phenolic content**

Sr. No.	Conc. µg/ml	Absorbance of Gallic acid (at 666 nm)
1.	0	0.000
2.	2	0.101 ± 0.0008
3.	4	0.141 ± 0.0008
4.	6	0.218 ± 0.0008
5.	8	0.307 ± 0.0008
6.	10	0.372 ± 0.0008

Each value represents the mean ± SEM (n-3), Gallic acid as a standard.



**Image No.1: Calibration curve & calibration equation of Gallic acid**




**Table 8** Total phenolic content of AR stem bark extracts

Sr. No.	Extract	Conc. µg/ml	Absorbance	TPC (mg/GAE/g DW)
1.	Ethyl acetate	50	0.203±0.003	107
2.	Methanol	50	0.175±0.002	92.8

Each value represents the mean ± SEM (n-3), Gallic acid as a standard.

Above observation table reveals that Ethyl acetate and methanol have Phenolic content as 107(mg GAE/g DW), 92.8 (mg GAE/g DW) respectively. Ethyl acetate extract shows more phenolic content than methanol as per comparative evaluation of phenolic content of extracts.

Note: mg/GAE/g DW denotes mg per Gallic Acid Equivalent per gram dry weight.



**Graph No.1:** TPC of *Annona reticulata* L.

## ii. Total Flavonoid Content

**Table 9:** Absorbance of Quercetin at different concentrations for determination of total Flavonoid content

Sr. No.	Conc. µg/ml	Absorbance of Quercetin (at 510 nm)
1.	0	0.000
2.	2	0.021 ± 0.0008
3.	4	0.033 ± 0.0008
4.	6	0.048 ± 0.0008
5.	8	0.072 ± 0.0008
6.	10	0.090 ± 0.0005





Image No.17: Calibration curve & calibration equation of Quercetin

Table 10 Total flavonoid content of AR Stem Bark extracts

Sr. No.	Extract	Conc. µg/ml	Absorbance	TFC (mg/QE/g DW)
1.	Ethyl acetate	50	0.071±0.0017	159.4
2.	Methanol	50	0.031±0.0029	69.6

Above observation table reveals that Ethyl acetate and methanol have Flavonoid content as 159.4 (mg/QE/g DW), 69.6 (mg/QE/g DW) respectively. Ethyl acetate extract shows more Flavonoid content than methanol as per comparative evaluation of flavonoid content of extracts. Note: mg/QE/g DW denotes mg per Quercetin Equivalent per gram dry weight.



Graph No.2: TFC of *Annona reticulata* L.


#### D. Pharmacological screening of Plant extracts for- *In-vitro* antioxidant activity

Table 11 Absorbance of Gallic acid, Rutin, Quercetin


Sr.No.	Conc. (µg/ml)	Absorbance		
		Gallic acid	Rutin	Quercetin
1	20	0.423 ± 0.002	0.251 ± 0.002	0.484 ± 0.02
2	40	0.380 ± 0.001	0.218 ± 0.001	0.375 ± 0.001
3	60	0.316 ± 0.002	0.193 ± 0.001	0.282 ± 0.001
4	80	0.287 ± 0.001	0.163 ± 0.002	0.194 ± 0.001
5	100	0.195 ± 0.002	0.150 ± 0.001	0.155 ± 0.002

**Table 12:** % of inhibition Gallic acid, Rutin, Quercetin

Sr.No.	Conc. µg/ml	% inhibition		
		Gallic acid	Rutin	Quercetin
1	20	48.30	69.26	44.4
2	40	53.59	73.33	54.20
3	60	61.37	76.43	65.56
4	80	64.95	80.09	76.22
5	100	76.14	81.68	88.92




**Graph No.3:** Absorbance of Quercetin + DPPH



**Graph No.4:** % of inhibition of Standard compounds

**Table 13** Absorbance of AR stem bark extracts for determination of Antioxidant activity

Sr. No.	Conc. (µg/ml)	Absorbance		%inhibition	
		Ethyl acetate extract	Methanol extract	Ethyl acetate extract	Methanol extract
1.	20	0.682±0.001	0.708±0.022	16.72	13.55
2.	40	0.491±0.003	0.656±0.0031	40.17	19.90
3.	60	0.391±0.003	0.580±0.004	52.25	29.18
4.	80	0.280±0.005	0.490±0.006	65.81	40.17
5.	100	0.167±0.008	0.391±0.007	79.60	52.25



**Graph No.5:** % of inhibition of AR stem bark extracts

In DPPH Scavenging assay method, the % inhibition of Ethyl acetate and methanolic stem bark extracts of *Annona reticulata* L. at 513 nm has been recorded at different concentration of 20, 40, 60, 80 and 100 µg/ml respectively. The results were compared with Quercetin as a reference standard and both extracts showed very significant % inhibition close to reference standard. Among both extracts, Ethyl acetate extract of *Annona reticulata* L. shows highest % inhibition activity (79.60% at 100µg/ml concentration).

#### IV. SUMMARY AND CONCLUSION

*Annona reticulata* L. stem bark extracts were used for studying Pharmacognostic, Phytochemical and Pharmacological evaluation. The present study aimed to evaluating the wound healing effect of crude extract of *Annona reticulata* L. stem bark on rat. Two extracts were prepared AR ethyl acetate and AR methanol.

Phytochemical study revealed that presence of Carbohydrates, Alkaloids, Glycosides, protein, phenol, flavonoid, steroids and tannin and was further confirmed by studying the Thin Layer Chromatography. *In vitro* antioxidant methods it is proved that *Annona reticulata* L. stem bark extract has a good antioxidant activity and may be useful in many diseases that occur due to oxidative stress.

The *in-vitro* antimicrobial study has showed that Ethyl acetate and Methanolic extract of *Annona reticulata* L. does possess significant antibacterial activity with 50 mg/ml concentrations and with high concentration of Ethyl acetate extract (100 mg/ml) it shows 8 mm zone of inhibition of microorganism *Staphylococcus aureus*. The Methanol extract shows zone of inhibition 5mm against microorganism *Staphylococcus aureus*.

The results obtained in present study indicate that AR Ethyl acetate and AR Methanol extracts have significant free radical scavenging activity.

The overall antioxidant activity of these extracts might be attributed to its flavonoids, phenolic, steroid and other phytochemical constituents. These could be source of natural antioxidant and have greater importance or showing oxidative stress related degenerative diseases.

## V. REFERENCES:

1. Ambiga, S., et al. "Evaluation of wound healing activity of flavonoids from Ipomoea carnea Jacq." *Ancient Science of Life* 26.3 (2007): 45.
2. Barku, Victor YA. et al. "Phytochemical screening and assessment of wound healing activity of the leaves of Anogeissus leiocarpus." *European Journal of Experimental Biology* 3.4 (2013): 18-25.
3. Barua, C. C. et al. "Evaluation of the wound healing activity of methanolic extract of Azadirachta Indica (Neem) and Tinospora cordifolia (Guduchi) in rats." *Pharmacologyonline* 1 (2010): 70-77.
4. Barua, C. C. et al. "wound healing activity of methanolic extract of leaves of Alternanthera brasiliana Kuntz using in vivo and in vitro model." *Indian journal of experimental biology* 47.12 (2009): 1001-1005.
5. Bhalke, Rasika Dnyandeo. et al. "Analgesic and CNS depressant activities of extracts of Annona reticulata Linn. bark." *Phytopharmacology* 1.5 (2011): 160-5.
6. Bharadwaj, R. et al. "Topical delivery of methanolic root extract of Annona reticulata against skin cancer." *South African Journal of Botany* 124 (2019): 484-493.
7. Biswas, Tuhin Kanti, et al. "Evaluation of Cynodon dactylon for wound healing activity." *Journal of ethnopharmacology* 197 (2017): 128-137.
8. Cai, Li. "Thin layer chromatography." *Current Protocols Essential Laboratory Techniques* 8.1 (2014): 6-3.
9. Chaudhari, Minakshi. et al. "Evaluation of phytoconstituents of Terminalia arjuna for wound healing activity in rats." *Phytotherapy Research: An International Journal Devoted to Pharmacological and Toxicological Evaluation of Natural Product Derivatives* 20.9 (2006): 799-805.
10. Chavan Akshay. et al. "Study of anthelmintic potential of ethanolic extract of Annona reticulata Linn leaves." *World journal of pharmacy and pharmaceutical sciences*. 9. (2020). 1352-1360.
11. Chavan, Machindra J. et al. "Analgesic and anti-inflammatory activities of the sesquiterpene fraction from Annona reticulata L. bark." *Natural product research* 26.16 (2012). 1515-1518.
12. Cobos, David Segura, et al. "Anti-inflammatory and antinociceptive properties of the extracts from the leaves of Porophyllum tagetoides and Annona reticulata." *Journal of Medicinal Plants* 7.1 (2019): 50-54.
13. Dash, G. K. et al. "Studies on Wound Healing Activity of Heliotropium indicum Linn. Leaves on Rats." *International Scholarly Research Network*. 2011. (2011) 1-8.
14. Ekaluo, U. B., et al. "In vitro antioxidant and free radical activity of some Nigerian medicinal plants: bitter leaf (Vernonia amygdalina L.) and guava (Psidium guajava Del.)." *Research Journal of Medicinal Plant* 9.5 (2015): 215-226.
15. Esimone, C. O. et al. "Cutaneous wound healing activity of a herbal ointment containing the leaf extract of Jatropha curcas L. (Euphorbiaceae)." *Int J Appl Res Nat Prod* 1.4 (2008): 1-4.
16. Fikru, Abraham, et al. "Evaluation of in vivo wound healing activity of methanol extract of Achyranthes aspera L." *Journal of ethnopharmacology* 143.2 (2012): 469-474.
17. Garg. et al. "Wound-healing activity of ethanolic and aqueous extracts of Ficus benghalensis." *Journal of advanced pharmaceutical technology & research* 2.2 (2011): 110.
18. Ghaisas. et al. "Evaluation of wound healing activity of ferulic acid in diabetic rats." *International wound journal* 11.5 (2014): 523-532.
19. Gulzar Alam. et al. "Wound healing potential of some medicinal plants." *International journal of pharmaceutical sciences review and research* 9.1. (2011): 36-145.
20. Hamde C. et al. "Formulation, evaluation and antimicrobial activity of Annona reticulata (ramphal) micro suspension." *World journal of pharmacy and pharmaceutical science* 9.10. (2020): 1841-1860.
21. Harish, B. G., et al. "Wound healing activity and docking of glycogen-synthase-kinase-3-beta-protein with isolated triterpenoid lupeol in rats." *Phytomedicine: International Journal of Phytotherapy and Phytopharmacology* 15.9 (2008): 763-767.
22. Islam, Md Rashedul, et al. "Antinociceptive activity studies with methanol extract of Annona reticulata L. (Annonaceae) and Carissa carandas L. (Apocynaceae) leaves in Swiss albino mice." *Adv. Nat. Appl. Sci* 6 (2012): 1313-1318.
23. Jahan, Sharmin, et al. "Antidiarrheal and antimotility activities of stem bark extracts of Annona reticulata Linn. in mice model." *J Appl Life Sci Int* 20.3 (2019): 1-9.
24. Jain Nilesh. et al. "Evaluation of wound-healing activity of Acorus calamus." *Natural product research*, 24:9 (2010): 534-541.
25. Jamkhande, Prasad G. et al. "Annona reticulata Linn. (Bullock's heart): Plant profile, phytochemistry and pharmacological properties." *Journal of Traditional and Complementary Medicine* 5.3 (2015): 144-152.
26. Kalyani Pathak. et al. "Comparative Pharmacological Evaluation on The Leaf and Stem Bark of Annona Reticulata L. for Antidiabetic Activity." *Nutraceutical & alternative medicine* (2013): 117-121.



27. Kalyani Pathak. et al. "Evaluation of leaf fraction of *Annona reticulata* for antioxidant activity and anti-inflammatory potential against carrageenan induced rat paw oedema model. " *Indian journal of natural sciences*,12,68. (2021).33877-3888.
28. Karodi, R., et al. "Evaluation of the wound healing activity of a crude extract of *Rubia cordifolia* L. (Indian madder) in mice. " *International Journal of Applied Research in Natural Products* 2.2 (2009): 12-18.
29. Katole, R. et al. "Phytochemical study of *Annona squamosa* L. and *Annona reticulata* L." *Int. J. Res* 5 (2018).156-175.
30. Khan, Nadeem. et al. "Unveiling antioxidant and anti-cancer potentials of characterized *Annona reticulata* leaf extract in 1, 2-dimethylhydrazine-induced colorectal cancer in Wistar rats." *Journal of Ayurveda and Integrative Medicine* 12.4 (2021): 579-589.
31. Khanam, Sofia. et al. "A systematic review on wound healing and its promising medicinal plants." *IP Int J Compr Adv Pharmacol* 5 (2021): 170-176.
32. Khandelwal K. R. et al. "Practical pharmacognosy" Ninth edition Nirali Prakashan Delhi, (2002) 25.1- 25.9.
33. Latha. S. et al. "Formulation and wound healing activity of seed extract of *Annona reticulata*. " *Indian drugs*,47:1. (2009): 48-51.
34. Mazumdar, Swagata, et al. "Immunomodulatory activity of ethanol extract of *Annona reticulata* L. leaf in cultured immune cells and in Swiss albino mice." *Journal of Ayurveda and Integrative Medicine* 13.2 (2022): 100554.